

# Project Overview: Impacts of Climate Change on the Phenology of Linked Agriculture-Water Systems

William P. Ball (1), Lisa Wainger (2a), Ciaran Harman (1), Damian Brady (3), Ariel Ortiz-Bobea (4), W. Michael Kemp (2b), Jeremy M. Testa (2a), Laura Murray (2b), Daniel Wilusz (1).


Johns Hopkins University, Dept. of Geography and Environmental Engineering; (2) University of Maryland Center for Environmental Sciences, (a) Chesapeake Bay Laboratory, (b) Horn Point Laboratory; (3) University of Maine, School of Marine Sciences; (4) Cornell University, School of Applied Economics and Management. **Contact:** Daniel Wilusz, dwilusz1@jhu.edu; William Ball, wball@jhu.edu

## Overarching Hypothesis

Climate-induced adaptive changes in agricultural practices and ecosystem processes will cause large indirect phenologic (seasonal and intra-annual) effects in riverine loading that will propagate through watersheds to the estuary and affect ecosystem health.

## Chesapeake Bay System Study Site

- Deep central channel with seasonal (summer) stratification.
- Large watershed to estuarine surface area ratio (~15:1).
- Increasing agricultural yields and development during the last 60 years.
- >50-years of monitoring data and >30-years of intense modeling effort in watershed and estuary; U.S EPA enforced TMDL activity since 2010.




**Figure 1.** The Chesapeake Bay watershed and major tributaries (left) and estuary (right).

## 1. Farmer Adaptations to Climate Change.

### How is farmer crop choice changing with climate?


- Discrete choice modeling using climatic, environmental, and economic factors.
- Analyzing climatic thresholds for double cropping systems.
- Developing methods for "big data" econometric analysis.
- Allowing for farm structural change.



**Figure 2.** Simulation of crop choice (upper panel) and timing of crop development (right panel).

### How are farmers altering timing of activities to manage risk?

- Crop yield modeling with varying sensitivities to weather shocks throughout growing season.
- Simulating alternative planting decisions (and subsequent fertilization timing).
- Combining timing and crop choice decisions to predict changes in nutrients available to run off.



## 2. Watershed Processes Affecting the Phenology of Connectivity.

**Motivation question**  
How do changes in climate and land use affect the timing and magnitude of water, N, P and SS loads reaching the Bay?

**Key science question**  
How do key, poorly understood watershed processes influence inter- and intra-annual transport dynamics?


### Lag times

- Using age-rank StorAge Selection (rSAS) functions to better represent lags in stream response to nutrient loading from groundwater.

### Variable source areas

- Using TOPOSWAT to simulate the influence of VSAs on inter-annual variability of nutrient transport.

**Figure 3.** Using transit time models to study groundwater transport lag as a function of changes in recharge. Includes data courtesy of Ward Sanford, USGS.




## 4. Adaptive Policy Instrument Evaluation.

### Identify policy-relevant leading indicators of environmental harm.

- Finding combinations of human actions and biophysical system responses that generate economic harms.
- Identifying the type, timing and scale of nutrient reduction interventions that avoid adverse conditions, e.g.:
  - Post-drought actions to capture unused nutrients
  - Payments for changes in tile drain management contingent upon rainfall patterns.

### Create rapid-response (state-contingent) policy tools to adaptively manage environmental risk.

- Computing the optimal interventions - where marginal cost of pollution reduction equals marginal damage cost of pollution delivered.
- Simulating adoption of policy instruments using risk-adjusted profit expectations and heterogeneous producers.
- Evaluating cost-effectiveness of dynamic instruments under likely adoption scenarios.




**Figure 5.** Considering policy costs and benefits.

## 3. Water Quality Outcomes in the Estuary.

### How do changes in the timing and location of watershed inputs affect estuary physical and biological responses, including the severity of hypoxia?

Analyzing 30-year Bay monitoring dataset for chlorophyll, dissolved oxygen, and nutrients to understand shifts in the seasonal cycles.

Conducting simulations using ROMS-RCA to understand the impacts of alternative loading seasonality and pulsed nutrient inputs on oxygen criteria failure, which will link to economics team.



**Figure 4.** Seasonal cycles of chlorophyll-a over each of the past three decades show changes in intra-annual variability (upper left panel). Simulated oxygen distribution in July and associated cumulative frequency diagram to assess criteria (upper right panel).

## Climate Change and Land Use (CC/LU) Scenarios

- Forcing scenarios with a common set of climate and land use scenarios representing plausible conditions up to 2100.
- Deriving climate scenarios from downscaled and bias-corrected CMIP global circulation models.
- Land use scenarios being made by partners using the Chesapeake Bay Land Change Model.

## Close Integration with Bay Management Activities

- Working closely with Chesapeake Bay Program Office (USEPA, USGS, NOAA) in support of Chesapeake Bay Program partnership.
- Helping improve the Chesapeake Bay Modeling System (toward 2017 and 2022 TMDL USEPA deadlines).
- Running workshops and symposia in collaboration with the Chesapeake Research Consortium.

## Educational Outreach

Hosting a school teacher to develop an educational module for the Teach Ocean Website. ([www.teachoceanscience.net](http://www.teachoceanscience.net)).

## Acknowledgements

We thank our project partners and collaborators. Primary project funding provided by four NSF WSC-3 grants for collaborative research: CBET-1360415, -1360345, -1360424, and -1360395.

